

REPORT DOCUMENTATION PAGE			2	Form Approved OMB NO. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA, 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 20-08-2014		2. REPORT TYPE Final Report		3. DATES COVERED (From - To) 1-Sep-2009 - 31-Aug-2013	
4. TITLE AND SUBTITLE Methods for Using Ab Initio Potential Energy Surfaces in Studies of Gas-Phase Reactions of Energetic Molecules			5a. CONTRACT NUMBER W911NF-09-1-0469		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER 611102		
6. AUTHORS Donald L Thompson			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAMES AND ADDRESSES University of Missouri - Columbia 310 Jesse Hall  Columbia, MO 65211 -1230			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS (ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			10. SPONSOR/MONITOR'S ACRONYM(S) ARO		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) 54142-CH.2		
12. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.					
14. ABSTRACT The focus of this research was to apply efficient methods for using ab initio potential energy surfaces (PESs) computed with high levels of quantum chemistry theory to predict chemical reaction properties of energetic molecules. The project has provided results of direct relevance to understanding the chemistry of energetic materials, particularly, 1,3,3-trinitroazetidine (TNAZ) decomposition.					
15. SUBJECT TERMS energetic materials, electronic structure, potential energy surfaces					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	15. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Donald Thompson
a. REPORT UU	b. ABSTRACT UU	c. THIS PAGE UU			19b. TELEPHONE NUMBER 573-882-0051

## Report Title

Methods for Using Ab Initio Potential Energy Surfaces in Studies of Gas-Phase Reactions of Energetic Molecules

### ABSTRACT

The focus of this research was to apply efficient methods for using ab initio potential energy surfaces (PESs) computed with high levels of quantum chemistry theory to predict chemical reaction properties of energetic molecules. The project has provided results of direct relevance to understanding the chemistry of energetic materials, particularly, 1,3,3-trinitroazetidine (TNAZ) decomposition.

---

**Enter List of papers submitted or published that acknowledge ARO support from the start of the project to the date of this printing. List the papers, including journal references, in the following categories:**

**(a) Papers published in peer-reviewed journals (N/A for none)**

Received

Paper

**TOTAL:**

**Number of Papers published in peer-reviewed journals:**

---

**(b) Papers published in non-peer-reviewed journals (N/A for none)**

Received

Paper

**TOTAL:**

**Number of Papers published in non peer-reviewed journals:**

---

### (c) Presentations

“A DFT and Ab Initio Study of the Thermal Decomposition of 1, 3, 3-Trinitroazetidine (TNAZ),” Jeffrey D. Veals and Donald L. Thompson, American Physical Society Prairie Section Fall 2013 Meeting, University of Missouri, Columbia, MO November 2013. (poster)

“A DFT and Ab Initio Study of the Thermal Decomposition of 1, 3, 3-Trinitroazetidine (TNAZ),” Jeffrey D. Veals and Donald L. Thompson, 40th National Organization for the Professional Advancement of Black Chemists and Chemical Engineers National Meeting, Indianapolis, IN October 2013. (poster)

---

**Non Peer-Reviewed Conference Proceeding publications (other than abstracts):**

Received      Paper

**TOTAL:**

---

**Number of Non Peer-Reviewed Conference Proceeding publications (other than abstracts):**

**Peer-Reviewed Conference Proceeding publications (other than abstracts):**

Received      Paper

**TOTAL:**

---

**Number of Peer-Reviewed Conference Proceeding publications (other than abstracts):**

---

**(d) Manuscripts**

Received      Paper

**TOTAL:**

---

**Number of Manuscripts:**

---

**Books**

Received      Book

**TOTAL:**

ReceivedBook Chapter**TOTAL:****Patents Submitted****Patents Awarded****Awards**

None

**Graduate Students**

<u>NAME</u>	<u>PERCENT SUPPORTED</u>	Discipline
Rezvan Chitsazi	0.69	
Jamin Perry	0.31	
Yi Shi	1.00	
Mehdi Shakourian Fardjahromi	0.31	
<b>FTE Equivalent:</b>	<b>2.31</b>	
<b>Total Number:</b>	<b>4</b>	

**Names of Post Doctorates**

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
Jeffrey Veals	0.50
<b>FTE Equivalent:</b>	<b>0.50</b>
<b>Total Number:</b>	<b>1</b>

**Names of Faculty Supported**

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
<b>FTE Equivalent:</b>	
<b>Total Number:</b>	

---

### **Names of Under Graduate students supported**

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
<b>FTE Equivalent:</b>	
<b>Total Number:</b>	

#### **Student Metrics**

This section only applies to graduating undergraduates supported by this agreement in this reporting period

The number of undergraduates funded by this agreement who graduated during this period: ..... 0.00

The number of undergraduates funded by this agreement who graduated during this period with a degree in science, mathematics, engineering, or technology fields:..... 0.00

The number of undergraduates funded by your agreement who graduated during this period and will continue to pursue a graduate or Ph.D. degree in science, mathematics, engineering, or technology fields:..... 0.00

Number of graduating undergraduates who achieved a 3.5 GPA to 4.0 (4.0 max scale):..... 0.00

Number of graduating undergraduates funded by a DoD funded Center of Excellence grant for Education, Research and Engineering:..... 0.00

The number of undergraduates funded by your agreement who graduated during this period and intend to work for the Department of Defense ..... 0.00

The number of undergraduates funded by your agreement who graduated during this period and will receive scholarships or fellowships for further studies in science, mathematics, engineering or technology fields:..... 0.00

---

### **Names of Personnel receiving masters degrees**

<u>NAME</u>
<b>Total Number:</b>

---

### **Names of personnel receiving PHDs**

<u>NAME</u>
<b>Total Number:</b>

---

### **Names of other research staff**

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
<b>FTE Equivalent:</b>	
<b>Total Number:</b>	

---

**Sub Contractors (DD882)**

**Inventions (DD882)**

**Scientific Progress**

See attachment.

## Technology Transfer

Several years ago we reported DFT calculations at the B3LYP/6-31g(d,p) level for the three main pathways commonly proposed for the decomposition mechanism of TNAZ (1,3,3-trinitroazetidine). [S. Alavi, L. M. Reily, and D. L. Thompson, *J. Chem. Phys.* **119**, 8297 (2003).] We considered the decomposition pathways in the mechanisms proposed by Zhang-Bauer [Y. X. Zhang and S. H. Bauer, *J. Phys. Chem. A* **102**, 5846 (1998).] and Lee. [D. S. Ames, J. C. Allman, Y. T. Lee, in *Chemistry of Energetic Materials*, G. A. Olah, D. R. Squire, Eds; (Academic Press: Sandiego, 1991) p. 27.] The common assumption is that the chemistry begins with either a C-N or the N-N bond fission. We reported values for the energies for N-NO<sub>2</sub> and C-NO<sub>2</sub> of 38.27 kcal/mol and 39.49 kcal/mol, respectively. We predicted the energy barriers for HONO elimination as 44.0 kcal/mol for formation from one of the C-NO<sub>2</sub> groups and 45.0 kcal/mol for formation from the N-NO<sub>2</sub> group.

Laura Butler (University of Chicago) brought to our attention a paper by Oftadeh and coworkers [M. Oftadeh, M. H. Khozani, M. Radhoosh, and M. H. Keshavarz, *Comp. Theo. Chem.*, **964**, 262 (2011).] in which the energy for C-N bond fission was reported to be 76.6 kcal/mol and 69.3 kcal/mol for the N-N bond rupture, and a 149.3 kcal/mol energy barrier for HONO elimination. This prompted a completely comprehensive investigation of the decomposition pathways using better density functionals than were available when we did our study in 2003. We performed geometry optimizations at the B3LYP level of theory with the 6-31G(d,p) Pople basis set, and the optimized structures were used as starting structures for M06 [Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.* **120**, 215 (2008).] and M06-2X**Error! Bookmark not defined.** geometry optimizations using the 6-31G(d,p) basis set. After these preliminary results involving the C-N and N-N initialization pathways we concluded that the M06 density functional should be used with the cc-pVTZ basis set to perform more accurate geometry optimizations for the three decomposition pathways and their products. This decision was based on the slightly better agreement between the M06 energies and the experimental values; we believe it is unlikely that our conclusions would have changed much if the M06-2X functional had been used for the optimizations instead.

To further improve the energies we performed M06/cc-pVQZ single point calculations on all the M06/cc-pVTZ optimized structures. Finally, CCSD(T)/cc-pVTZ energies were obtained using the M06/cc-pVTZ geometries. The *ab initio* calculations were done as a precaution as well as another a means of gaining insight about the initial decomposition process. This extra effort was made in large part to shore up any final conclusions to be drawn our study.


We performed the M06/cc-pVTZ optimizations and CCSD(T)/cc-pVTZ energies for the decomposition steps subsequent to these initial reactions because for several cases, particularly ring opening, a multi-configuration-based method would be needed to gain any further clarity. Since M06 was optimized for organometallics as well as typical organic molecules, we believe it is probably more accurate than other DFT (or MP2) methods for somewhat multi-configurational (bi-radicals, highly resonant, etc) species, and that only a dedicated multi-reference method would significantly improve upon the M06 predicted geometries, we settled for the latter. We are fairly confident that the M06 results give good qualitative descriptions of likely decomposition routes.

This work has been published: Jeffrey D. Veals and Donald L. Thompson, "Thermal Decomposition of 1,3,3-Trinitroazetidine (TNAZ): A Density Functional Theory and *Ab Initio* Study," *J. Chem. Phys.* **140**, 154306 (2014). The abstract of this paper is:

Density functional theory and ab initio methods are employed to investigate decomposition pathways of 1,3,3-trinitroazetidine initiated by unimolecular loss of  $\text{NO}_2$  or HONO. Geometry optimizations are performed using M06/cc-pVTZ and coupled-cluster (CC) theory with single, double, and perturbative triple excitations, CCSD(T), is used to calculate accurate single-point energies for those geometries. The CCSD(T)/cc-pVTZ energies for  $\text{NO}_2$  elimination by N-N and C-N bond fission are, including zero-point energy (ZPE) corrections, 43.21 kcal/mol and 50.46 kcal/mol, respectively. The decomposition initiated by trans-HONO elimination can occur by a concerted H-atom and nitramine  $\text{NO}_2$  group elimination or by a concerted H-atom and nitroalkyl  $\text{NO}_2$  group elimination via barriers (at the CCSD(T)/cc-pVTZ level with ZPE corrections) of 47.00 kcal/mol and 48.27 kcal/mol, respectively. Thus, at the CCSD(T)/cc-pVTZ level, the ordering of these four decomposition steps from energetically most favored to least favored is:  $\text{NO}_2$  elimination by N-N bond fission, HONO elimination involving the nitramine  $\text{NO}_2$  group, HONO elimination involving a nitroalkyl  $\text{NO}_2$  group, and finally  $\text{NO}_2$  elimination by C-N bond fission.


Below are figures that summarize our results

**Energy profile for decomposition beginning with N-N bond fission:**





Potential energy profile for decomposition beginning with C-N bond fission:



Potential energy profile for HONO elimination:

